Anti-Calreticulin / CALR antibody, rabbit polyclonal

Product code	73-707
Size	100 µg
Storage	-20℃
Concentration	1mg/ml
Buffer	PBS- with 50 % glycerol
Purity	Purified IgG fraction with protein A from rabbit antiserum (73-018).
Immunogen	C-EEDEKEEDEEESPGQAKDEL (C-terminal of mouse CALR protein)
Isotype	Rabbit IgG
Reactivity	Mouse and human. Not tested with other species.
Special notes	Validation: Knock-out mouse
Application	1. Western blotting (1/500~1/1,000 dilution) 2. Immunoprecipitation (1/100) 3. Immunoflourescence staining (1/300~1/1,000 dilution) 4. Immunohistochemistry (1/1,000 dilution) Other applications have not been tested.
Background	Calreticulin (CALR) is calcium-binding chaperone that promotes folding, oligomeric assembly and quality control in the endoplasmic reticulum (ER) via the calreticulin/calnexin cycle. This lectin interacts transiently with almost all of the monoglucosylated glycoproteins that are synthesized in the ER. Interacts with the DNA-binding domain of NR3C1 and mediates its nuclear export. Involved in maternal gene expression regulation. May participate in oocyte maturation via the regulation of calcium homeostasis
Data Link	uniprot/P14211 Mouse Calrecticulin, uniprot/Q96L12 Human Calreticulin Gene I D 12317 Mouse Calr, Gene ID 811 Human Calr
Reference	This antibody was described and used in the following publication. 1. Ikawa M. et al (2011) Calsperin is a testis-specific chaperone required for sperm fertility. J Biol Chem.18:5639-46. pubmed/21131354 Free article. **WB, IP**
Related Products	73-018 Anti-Calreticulin / CALR antibody, rabbit serum

Please note: All products are FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR MILITARY USE.

BioAcademia,Inc. Tel. 81-6-6877-2335 Fax. 81-6-6877-2336 info@bioacademia.co.jp https://www.bioacademia.co.jp/en/

Data Images: 73-707 Anti-Calreticulin / CALR antibody, rabbit polyclonal




Fig.1 Identification of CALR protein by western blotting with anti-CALR antibody (73-018).

Embryonic fibroblast cells prepared from *Calr* -/- mouse were transfected with a plasmid expressing *Calr*. The cell lysate was analyzed by western blotting with anti-CALR antibody at 1/500 dilution.

1. Mock-infected cell lysate as a negative control.
2. Cell lysate transfected with a plasmid expressing *Calr*.

The molecular mass is 48 kDa




Fig.2 Western blot analysis of CALR protein in crude extracts of mouse testis and sperm with anti-CALR antibody (73-018).

Proteins in the extracts (10 µg protein) were separated on 10-20% gradient gel of SDS-PAGE and electro-blotted to a PVDF membrane. The membrane was reacted with anti-CALR antibody at 1/1,000 dilution. As the 2nd antibody, anti-rat IgG antibody conjugated with HRP (ab97051) was used at 1/10,000 dilution

BioAcademia,Inc. Tel. 81-6-6877-2335 Fax. 81-6-6877-2336 info@bioacademia.co.jp https://www.bioacademia.co.jp/en/




Fig.3 Western blot analysis of CALR protein in crude extracts of human cell lines with anti-CALR antibody (73-018, 73-707).

Proteins in the extracts (10 μg protein) were separated on 12.5% gel of SDS-PAGE and electro-blotted to a PVDF membrane (wet system). The membrane was reacted with anti-CALR antibody at 1/1,000 dilution. As the 2nd antibody, anti-rat IgG antibody conjugated with HRP (ab97051) was used at 1/10,000 dilution. Lane1. HeLa cell extracts Lane2. MCF-7 cell extracts.




Fig.4. Immunoprecipitation of CALR protein with anti-CALR antibody (73-018).

Lysates of wild-type mouse testis were immunoprecipitated with anti-CALR antibody and the precipitates were analyzed by western blotting with the same antibody.

1. Input testis lysate
2. Precipitated with preimmune serum
3. Precipitated with anti-CALR antibody

BioAcademia,Inc. Tel. 81-6-6877-2335 Fax. 81-6-6877-2336 info@bioacademia.co.jp https://www.bioacademia.co.jp/en/




Fig.5　Immunofluorescence staining of CALR protein in NIH3T3 cells with anti-CALR antibody (73-018).

NIH3T3 cells were fixed with 4% paraformaldehyde and permeabilized with 0.5% TritonX 100 and reacted with anti-CALR antibody at 1/300 dilution. As the 2nd antibody, goat anti-rabbit IgG antibody conjugated with Alexa Fluor 488 (Molecular Probes) was used at 1/1,000 dilution.
DNA was stained with DAPI (1 ug/ml) and the merged image was shown on right.




Fig.6. Immunohistological staining of CALR protein in mouse testis using anti-CALR antibody (73-018).

A section of formalin fixed and paraffin embedded mouse testis was treated with the anti-CALR antibody at 1/1,000 dilution after deparaffization and antigen retrieval. The 2nd antibody, goat anti-rabbit IgG conjugated with Alexa Fluor 488 (Molecular Probes #1166843) was used at 1/1,000 dilution. DNA was stained with DAPI (1.0μg/mL) and the merged image was shown (Merge). The bright-field image of the same region was shown on the right.

BioAcademia,Inc. Tel. 81-6-6877-2335　Fax. 81-6-6877-2336 info@bioacademia.co.jp　https://www.bioacademia.co.jp/en/